


What Is Claimed Is:

1. A method for immobilizing nucleic acid on a solid phase substrate comprising:

bringing the solid phase substrate into contact with a
5 composition comprising a total concentration of 0.1 to 2 μM of
a nucleic acid as a probe and a compound or a salt thereof,
the compound being represented by the following formula:




wherein L¹ is a single bond or a C₁₋₁₅ alkylene group; L² is a
10 single bond, a nucleic acid, a polyethylene glycol group, -CO-
NH-, or -NH-CO-; R is a hydroxyl group, an amino group, a
ferrocenyl group, or a carboxyl group; provided that neither L¹
nor L² is a single bond, and

incubating the composition in contact with a surface of
15 the solid phase substrate.


2. The method according to claim 1, wherein the nucleic acid as a probe comprises a polynucleotide or an oligonucleotide consisting of modified or unmodified, single-stranded DNA, RNA, PNA, CNA, or HNA.

20 3. The method according to claim 1, wherein the nucleic acid as the probe comprises at the 3' end or the 5' end a group represented by the following formula:



wherein L³ is a C₁₋₁₅ alkylene group, and L⁴ is a single
25 bond or a spacer.

4. The method according to claim 1, wherein the nucleic acid as the probe has at the 5' end a group represented by the following formula:



30 wherein L^4 is a single bond or a spacer.

5. The method according to claim 4, wherein L⁴ is a nucleic acid, -CO-NH-, -NH-CO-, a polyethylene glycol group, or a polyethylene glycol phosphate group.

6. The method according to claim 1, wherein the total concentration of the nucleic acid and the compound I or the salt thereof in the composition is 0.5 to 1.5 μM .

7. The method according to claim 1, wherein the total
5 concentration of the nucleic acid and the compound I or the salt thereof in the composition is 1 μM .

8. The method according to claim 1, wherein the composition comprises the nucleic acid and the compound I at a ratio of 40/60 to 60/40.

10 9. The method according to claim 1, wherein R in the formula I is a hydroxyl group.

10. The method according to claim 1, wherein L^1 in the formula I is a single bond, and L^2 is a polyethylene glycol group.

15 11. The method according to claim 1, wherein L^1 in the formula I is a C_{4-8} alkylene group, and L^2 is a single bond.

12. The method according to claim 1, wherein the compound I is 6-mercapto-1-hexanol.

13. The method according to claim 1, wherein the solid
20 phase substrate is a single layered substrate or a multiple layered substrate comprising at least one material selected from the group consisting of glass, polymer resin and metal.

14. The method according to claim 1, wherein the surface
25 of the solid phase substrate on which nucleic acid is adsorbed is coated with a thin gold film.

15. The method according to claim 1, wherein the solid phase substrate is a glass substrate with a thin gold film vapor-deposited on its surface, and may further comprises, at least one intermediate layer between the thin gold film and
30 the glass substrate.

16. The method according to claim 1, wherein the nucleic acid as the probe has 15 to 30 base length.


17. The method according to claim 1, wherein the incubation is carried out at a temperature of 25° C to 40° C.

18. The method according to claim 1, wherein the nucleic acid as the probe is a polynucleotide or an oligonucleotide
5 consisting of single-stranded DNA, RNA, or PNA, and may also has the group represented by formula II, the compound I is 6-mercapto-1-hexanol; the total concentration of the nucleic acid and 6-mercapto-1-hexanol in the composition is 0.5 to 1.5 μ M; and the solid phase substrate is a glass substrate with a
10 thin gold film vapor-deposited on its surface, and further, at least one intermediate layer may be made between the thin gold film and the glass substrate.

19. A method for manufacturing a biosensor having a nucleic acid probe as a sensing site comprising the use of the
15 nucleic acid immobilization method according to any of claims 1 to 18.

20. A biosensor comprising a solid phase substrate and a nucleic acid probe thereon as a sensing site, the biosensor manufactured by a method comprising the steps of:

20 bringing a composition containing a nucleic acid and a compound or a salt thereof at a total concentration of 0.5 to 1.5 μ M into contact with the solid phase substrate, the compound being represented by the following formula:



25 wherein L^1 is a single bond or a C_{1-15} alkylene group; L^2 is a single bond, a nucleic acid, a polyethylene glycol group, $-\text{CO}-\text{NH}-$, or $-\text{NH}-\text{CO}-$; R is a hydroxyl group, an amino group, a ferrocenyl group, or a carboxyl group; provided that neither L^1 nor L^2 is a single bond, and

30 incubating the composition in contact with a surface of a solid phase substrate.

21. The biosensor according to claim 20, wherein the compound I is 6-mercapto-1-hexanol, the solid phase substrate

is a glass substrate with a thin gold film vapor-deposited on its surface, and further, the solid phase substrate may comprises at least one intermediate layer between the thin gold film and the glass substrate.

5 22. A method for detecting a target nucleic acid molecule in a test sample by detecting a measurable signal, comprising the steps of:

 (a) bringing the test sample into contact with a biosensor having a nucleic acid probe manufactured by using
10 the nucleic acid immobilization method according to claim 1, and

 incubating the test sample in contact with a surface of the biosensor;

 (b) applying light to the solid phase substrate of the
15 biosensor from the opposite side of the surface to which nucleic acid is immobilized, continuously or intermittently from before to after step (a); and

 (c) measuring a shift of reflectivity of the solid phase substrate by detecting the reflection of the light applied in
20 step (b).

 23. The method according to claim 22, wherein the method is used for detecting DNA, RNA, or single nucleotide polymorphisms.

 24. A use of a composition comprising a nucleic acid and
25 compound I or salt thereof at a total concentration of 0.5 to 1.5 μM , for manufacturing a biosensor having a nucleic acid probe as a sensing site.

 25. A use of a nucleic acid and compound I or salt thereof, for manufacturing the composition for the use
30 according to claim 24.